



Science of Nursing
and Health Practices



Science infirmière
et pratiques en santé

Article de protocole de recherche | Research protocol article

The Different Types of Overtime Work in Nursing and Their Associations With Nurse and Patient Outcomes: A Cross-Sectional Study Protocol

Les différents types d'heures supplémentaires et leurs associations avec les résultats chez l'infirmière et le patient: un protocole d'étude transversale

Raouaa Braiki  <https://orcid.org/0000-0002-7765-9589> School of Nursing, Université de Sherbrooke, Quebec, Canada

Christian Rochefort  <https://orcid.org/0000-0002-0035-8382> School of Nursing, Université de Sherbrooke, Centre hospitalier universitaire de Sherbrooke, Centre de recherche de l'Hôpital Charles-Le Moyne, Quebec, Canada

Correspondance | Correspondence:

Raouaa Braiki

raouaa.braiki@usherbrooke.ca



2025 R Braiki, C Rochefort.

[Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

ISSN 2561-7516

Keywords	Abstract
overtime; mandatory; voluntary; nurse; quality and safety of care	<p>Introduction: Nurses often work overtime to fill the shortage of nurses, ensure continuity of care or prevent service breakdowns. Some studies show that working overtime has negative impacts on both nurse and patient outcomes, whereas others suggest that overtime has some beneficial outcomes for patients and nurses. Some authors suggest that these conflicting results across studies could be explained by the type of overtime performed by nurses, an aspect that has received scant research attention. Objective: We aim to examine the associations between the different types of overtime work (voluntary or mandatory), and nurses' perceptions of nurse and patient outcomes. Method: A provincial electronic cross-sectional survey will be conducted in the province of Quebec, Canada, to examine the associations between nurse overtime work and both nurse and patient outcomes. Discussion and Research Spin-offs: This study will likely provide deeper insights about the different types of overtime and their impacts on both nurse and patient outcomes. This may inform nursing practices and guide nursing union representatives, directors of nursing and government decision-makers.</p>

Résumé	Mots-clés
<p>Introduction : Les infirmières travaillent souvent des heures supplémentaires pour combler la pénurie d'infirmières, assurer la continuité des soins ou pour prévenir des bris de services. Plusieurs études suggèrent qu'une augmentation du travail en heures supplémentaires est associée à de moins bons indicateurs de la santé au travail des infirmières, ainsi qu'à des soins de moindre qualité et moins sécuritaires; d'autres n'ont observé aucune association significative, ou encore un effet protecteur des heures supplémentaires. Certains auteurs suggèrent que ces résultats contradictoires pourraient s'expliquer par le type d'heures supplémentaires effectuées par les infirmières, un aspect qui a reçu peu d'attention dans les recherches. Objectif : Examiner les associations entre les différents types d'heures supplémentaires (volontaires ou obligatoires) et les perceptions des infirmières à l'égard des résultats chez les infirmières et les patients. Méthode : Une étude transversale sera réalisée auprès des infirmières travaillant dans les hôpitaux de la province de Québec, au Canada, pour examiner les associations entre les heures supplémentaires et les résultats des infirmières et des patients. Discussion et retombées anticipées : Cette étude fournira des connaissances plus approfondies sur les différents types d'heures supplémentaires et leurs conséquences sur l'infirmière et le patient, ce qui pourrait informer la pratique infirmière et guider les décisions des représentants des syndicats infirmiers, les gestionnaires et les décideurs gouvernementaux.</p>	heures supplémentaires; obligatoire; volontaire; infirmière; qualité et sécurité des soins

INTRODUCTION

Overtime work is defined as working more than the standard working hours (Anxo & Karlsson, 2019; Government of Canada, 2024). The excessive use of overtime to manage the nursing shortage and the variability in patient needs is a global issue in terms of its impacts on patient and nurse outcomes, as well as on health care system costs (Bae et al., 2014). In Canada, for example, nurses worked more than 15.2 million hours of paid overtime in 2016, which was then estimated to cost CDN \$788 million (Jacobson Consulting Inc., 2017). In 2022, Canadian nurses reported an average of 8.2 overtime hours per week (Statistics Canada, 2023). In the United States, among more than 5,000 registered nurses (RNs) surveyed by Medscape (2020); 56% were working between 1 and 5 hours of overtime per week, which cost USD 9,000 in additional annual wages per nurse (Medscape, 2020).

Numerous studies show that working overtime has negative impacts on patient and nurse outcomes (Bae et al., 2024; Cha et al., 2023; Gao et al., 2023; Min et al., 2019). According to the recent findings from the Canadian Community Health Survey, working overtime decreases job satisfaction and job security for nurses, and increases work stress (48.5% reported high work stress) (Statistics Canada, 2020). Also, working overtime was associated with career choice regret (Dyrbye et al., 2020); intent to leave (Bae et al.); burnout (Cha et al.; Chen et al., 2022; Gao et al.); fatigue (Min et al.) and poor quality of sleep (Caruso et al., 2019), which might decrease vigilance and alter decision-making ability (Geiger-Brown, Tinkoff, & Rogers, 2011; Trinkoff et al., 2011). In addition, working overtime decreases the quality of nursing service (Ball et al., 2017) and increases missed nursing care (Min et al.), and the occurrence of adverse patient outcomes (e.g., medication errors (Kunaviktiluk et al., 2015), infections (Beltempo et al., 2017; Stone et al., 2017), and patient mortality (Trinkoff et al.)). However, some studies found no significant or negative associations between working overtime and nurse outcomes (Bannai & Tamakoshi, 2014; Watanabe, Imamura, & Kawakami, 2016) or

patient outcomes (Berney & Needleman, 2006; Stone, 2007).

There are two main types of overtime work: Voluntary overtime (when a nurse consents to working extra hours to meet daily patient requirements for nursing care) and mandatory overtime (when a nurse is forced to work overtime by her/his employer to avoid a break in service). As such, some authors have proposed that the inconsistencies across studies could be explained by the type of overtime performed by nurses (Bougie & Cara, 2008; de Castro et al., 2010; Nogues & Tremblay, 2023; Watanabe & Yamauchi, 2016; 2018; 2019). However, to the best of our knowledge, this hypothesis has received scant attention in the literature.

OBJECTIVE

The objective of this study is to examine the associations between the type of overtime work (voluntary or mandatory), and nurses' perceptions of nurse and patient outcomes.

METHOD

STUDY DESIGN

This paper is a cross-sectional study protocol and is reported according to ObsQual checklist (Low et al., 2024). Thus, using a provincial electronic survey, a cross-sectional study will be performed. We will report this study according to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) criteria (von Elm et al., 2007).

POPULATION, SAMPLE AND SETTING

The study population consists of all RNs providing direct patient care in a medical, surgical, intensive care, or geriatric care unit in an acute care hospital in the province of Quebec, Canada ($N = 37\,819$). To ensure adequate statistical power for conducting multiple regression analyses (described below), we used the G*Power software (Faul et al., 2009) to estimate the required sample size. The calculation was based on the following parameters: A small effect size ($f^2 = 0.02$), a two-tailed alpha level of 0.05, a power of 0.80 and 17

predictors (including voluntary overtime, mandatory overtime and confounders). Based on these parameters, the required sample size is estimated to range from 6,667 to 10,000 participants, assuming an expected response rate of 10-15%. This range represents the final targeted sample size for the study.

Nurses will be recruited through two steps. First, we will ask the Quebec's Board of Nurses (*Ordre des infirmières et infirmiers du Québec*) to provide us with a random sample of 10 000 nurses, based on their list of members. Second, we will follow Dillman's et al. (2014) recommendations for electronic surveys. Specifically, an initial email invitation to potential study participants will be sent (T_0). Non-responders will receive reminders on day 4, 7 and 14 (Dillman et al.). An online survey link including a cover page and an electronic consent form will be sent to each potential participant.

DATA COLLECTION

We will use the RedCap platform to build and administer the electronic survey. Data collection will occur during winter 2025 through an electronic questionnaire. This questionnaire will contain the following sections: 1) overtime work characteristics; 2) process outcomes; 3) nurse job outcomes; 4) patient outcomes; and 5) confounders.

MEASURES

1) Overtime Work (Independent Variable)

It will be defined as "any hours worked in excess of the standard hours" (Government of Canada, 2024). Nurses will be asked to report on two overtime-related variables: a) the number of voluntary overtime hours performed over the previous 2 weeks (i.e., over the previous pay period); and b) the number of mandatory overtime hours performed over the previous 2 weeks.

2) Nurse Job Outcomes-Instruments

Nurses' perceptions on the following job outcomes will be measured:

- Job satisfaction will be measured using a single item extracted from the French version (Dupret et al., 2012) of the Copenhagen Psychosocial Questionnaire

Scale (COPSOQ) (International COPSOQ Network, 2025). The Job satisfaction of professional experience item will be scored on a 4-point Likert-type scale ranging from 1 (very satisfied) to 4 (very dissatisfied).

- Turnover intentions will be assessed using two items adapted from O'Driscoll & Beehr (1994): a) "I am thinking about leaving my job within the next 12 months" (Fernet et al., 2017); b) "I plan on searching for a new job outside the nursing profession within the next 12 months" (Lavoie-Tremblay et al., 2014). Items will be scored on a 7-point scale ranging from 1 (strongly disagree) to 7 (strongly agree).
- Emotional exhaustion will be measured using the emotional exhaustion subscale of the French version of the Maslach Burnout Inventory-Human Services Survey (MBI-HSS) (Dion & Tessier, 1994). Based on previous studies, emotional exhaustion is the most useful dimension of MBI-HSS (Aiken et al., 2023; White et al., 2019). This dimension will be assessed using 9 items. Participants will be asked to rate the frequency of experiencing feelings related to emotional exhaustion by one's work (e.g., "I feel depressed at work"); using a 7-point scale ranging from 0 (never) to 6 (daily). The emotional exhaustion subscale has excellent internal consistency (Lheureux et al., 2017).
- Fatigue will be measured using the French version of the Occupational Fatigue Exhaustion/Recovery (OFER-15) scale (Winwood et al., 2005). The OFER-15 is composed of three subscales: a) Acute Fatigue; b) Chronic Fatigue/Exhaustion; and c) Intershift recovery comprising 5 items each. Items will be scored on a 7-point Likert scale ranging from 0 (strongly disagree) to 6 (strongly agree). The OFER-15 has high internal consistency, with Cronbach alphas ranging from 0.84 to 0.86 across subscales. Also, temporal stability, construct and discriminant validity were acceptable (Winwood et al.).

3) Process Instrument

Missed nursing care is defined as care that is wholly or partially uncompleted or delayed (Kalisch & Williams, 2009). It will be measured using the MISSCARE Survey instrument (Kalisch & Williams), one of the most widely used tools in international research (Palese et al., 2020). Prior to administration, the instrument will be translated for the French-Canadian context following established translation guidelines; this process will be detailed elsewhere. The MISSCARE Survey instrument consists of two parts: A (elements of missed nursing care = 24 items) and B (reasons for missed nursing care = 17 items). As the study aims to examine the association between overtime and missed care, only part-A will be used. Participants will be asked to rate the frequency of each nursing care element on their unit over the past 2 weeks. Consistent with the original version, responses will be recorded using a 4-point Likert-type scale ranging from 1 (rarely missed) to 4 (always missed) (Kalisch & Williams). The instrument has demonstrated good internal consistency with Cronbach's alpha coefficient varying from 0.64 to 0.86 (Kalisch & Williams).

4) Patient Outcomes-Instruments

Nurses' perceptions of the frequency and quality of the following patient outcomes will be measured:

- Quality of care: Perceived quality of care on the nurses' unit over the past 14 days will be assessed using a single item of the French version (Rochefort, 2009) of the questionnaire developed by Aiken et al. (2001). Responses will be rated on a 5-point scale ranging from "poor" to "excellent".
- Safety of care: Perceived safety of care in the nurses' unit over the past 14 days will be measured by a single item from the French version (Occelli et al., 2013) of the Hospital Survey on Patient Safety Culture (Sorra & Dyer, 2010). Responses will be scored on a 5-point scale ranging from "failing" to "excellent".

CONFFOUNDERS

Potential confounding variables that may influence the relationship between overtime and

missed nursing care or patient outcomes will be measured. These include both **nurse-level** and **hospital-level** characteristics.

Nurses' characteristics will be assessed using 12 items: a) sex, b) gender, c) age, d) number of years of experience, e) level of education (college or bachelor's degree and above), f) marital status, g) number of children, h) personal income, i) subjective workload during the last 2 weeks, j) work position (full-time or part-time), k) work shift (day, evening, night, rotational), l) shift length (8 or 12 hours).

Hospital characteristics will be assessed using two items: a) type of hospital (teaching hospital VS. community hospital) and b) type of nursing unit.

ANALYSIS

Data analysis will be conducted with SPSS 26 (IBM, 2019). First, outliers and atypical responses will be removed from the dataset following the guidelines of Tabachnick & Fidell (2019). Missing data will be treated using multiple imputation methods, as recommended by Azur et al. (2011).

Descriptive analysis will be used to summarize participants' socio-demographic and hospital characteristics (e.g., mean, standard deviation); and to estimate the prevalence of each type of overtime. Then, correlation matrices will be computed among independent, control and outcome variables to assess potential multicollinearity and to inform the selection of variables for inclusion in subsequent linear regression models. Multiple linear regression models will be used to examine associations between each type of overtime (mandatory and voluntary) and selected nurse and patient outcomes. The assumptions of linear regression will be verified both a priori, via descriptive statistics and graphical inspection, and a posteriori, through analysis of regression residuals. To account for the clustering of nurses within units and hospitals, a Generalised Estimating Equations framework will be applied. The level of statistical significance will be set at $\alpha \leq 0.05$ with adjustments for multiple comparisons where applicable.

ETHICAL CONSIDERATIONS

This study was reviewed and approved by the research ethics committee of the *Université de*

Sherbrooke (no 2024-4590). Electronic consent will be available to participants on the RedCap platform. A cover page and an electronic information form will explain the study objective, eligibility criteria, expected participation and the principles of anonymity, confidentiality, and the right to accept, refuse or withdraw at any time. The questionnaires will be anonymous. We will assign a code for each questionnaire. Data will be stored for 10 years in a password-protected computer. Before conducting the provincial electronic survey, the adapted instrument (MISSCARE Survey) will be submitted to the research ethics committee for approval.

DISCUSSION AND RESEARCH SPIN-OFFS

This study aims to explore the characteristics of nurse overtime and to examine the associations of voluntary and mandatory overtime with nurse and patient outcomes. Findings from this study may provide evidence for the prevention of adverse outcomes of voluntary and mandatory overtime on both nurse and patient outcomes.

To our knowledge, this study will be the first to explore this topic in the province of Quebec, which has the highest overtime rates in Canada (Fédération interprofessionnelle de la santé du Québec, 2017). Furthermore, this study will investigate the perceptions of nurses working in several hospitals in the province of Quebec, which will provide deeper information about the reality of voluntary and mandatory overtime across the province. It will enable broad descriptions and comparisons. Another strength is the use of international and validated instruments. However, this study may have some limitations. First, the cross-sectional design of this study limits causal inferences. Second, the use of self-reported measures may lead to biases (e.g., recall bias, social desirability bias and common source), which could affect the study findings. Third, this study will not be looking at some confounders (e.g., patient ratio, staffing in units, and outbreaks) that may influence the association between working overtime (mandatory and voluntary) and patient and nurse outcomes.

Regarding our dissemination plan, different methods will be used to provide the maximum visibility for our project's results. First, we will publish the study project protocol in a scientific journal. Then, study findings will be disseminated in conferences and in at least four articles published in peer-reviewed journals. We will also seek opportunities to present our results to various stakeholders, including nursing union representatives, directors of nursing and other decision-makers.

Authors' contribution: RB and CR designed the study. RB wrote the first draft of this cross-sectional study protocol. CR revised and approved the final version of the manuscript.

Acknowledgments: None.

Funding: The authors received no funding to conduct the project reported in this article nor to draft this article.

Statement of conflict of interest: The authors declare no conflict of interest.

Reçu/Received: 5 Déc/Dec 2024 **Publié/Published:** 20 Août/August 2025

REFERENCES

Aiken, L. H., Lasater, K. B., Sloane, D. M., Pogue, C. A., Fitzpatrick Rosenbaum, K. E., Muir, K. J., McHugh, M. D., & US Clinician Wellbeing Study Consortium (2023). Physician and Nurse Well-Being and Preferred Interventions to Address Burnout in Hospital Practice: Factors Associated With Turnover, Outcomes, and Patient Safety. *JAMA health forum*, 4(7), e231809. <https://doi.org/10.1001/jamahealthforum.2023.1809>

Anxo, D., & Karlsson, M. (2019). *Overtime work: A review of literature and initial empirical analysis* (Conditions of work and employment series no. 104). International Labour Office. https://www.ilo.org/sites/default/files/wcmsp5/groups/public/%40ed_protect/%40protrav/%40travail/documents/publication/wcms_663072.pdf

Azur, M. J., Stuart, E. A., Frangakis, C., & Leaf, P. J. (2011). Multiple imputation by chained equations: what is it and how does it work?. *International journal of methods in psychiatric research*, 20(1), 40–49. <https://doi.org/10.1002/mpr.329>

Bae S. H. (2024). Nurse Staffing, Work Hours, Mandatory Overtime, and Turnover in Acute Care Hospitals Affect Nurse Job Satisfaction, Intent to Leave, and Burnout: A Cross-Sectional Study. *International journal of public health*, 69, 1607068. <https://doi.org/10.3389/ijph.2024.1607068>

Bae, S. H., & Fabry, D. (2014). Assessing the relationships between nurse work hours/overtime and nurse and patient outcomes: systematic literature review. *Nursing outlook*, 62(2), 138–156. <https://doi.org/10.1016/j.outlook.2013.10.009>

Ball, J., Day, T., Murrells, T., Dall’Ora, C., Rafferty, A. M., Griffiths, P., & Maben, J. (2017). Cross-sectional examination of the association between shift length and hospital nurses job satisfaction and nurse reported quality measures. *BMC Nursing*, 16, 26. <https://doi.org/10.1186/s12912-017-0221-7>

Bannai, A., & Tamakoshi, A. (2014). The association between long working hours and health: a systematic review of epidemiological evidence. *Scandinavian journal of work, environment & health*, 40(1), 5–18. <https://doi.org/10.5271/sjweh.3388>

Beltempo, M., Blais, R., Lacroix, G., Cabot, M., & Piedboeuf, B. (2017). Association of Nursing Overtime, Nurse Staffing, and Unit Occupancy with Health Care-Associated Infections in the NICU. *American Journal of Perinatology*, 34(10), 996–1002. <https://doi.org/10.1055/s-0037-1601459>

Berney, B., & Needleman, J. (2006). Impact of nursing overtime on nurse-sensitive patient outcomes in New York hospitals, 1995–2000. *Policy, politics & nursing practice*, 7(2), 87–100. <https://doi.org/10.1177/1527154406291132>

Bougie, M., & Cara, C. (2008). Temps supplémentaire obligatoire. Une première étude phénoménologique sur les perceptions d’infirmières. Sept thèmes se dégagent de leurs propos. *Perspective infirmière*, 5(8), 32–40.

Caruso, C. C., Baldwin, C. M., Berger, A., Chasens, E. R., Edmonson, J. C., Gobel, B. H., Landis, C. A., Patrician, P. A., Redeker, N. S., Scott, L. D., Todero, C., Trinkoff, A., & Tucker, S. (2019). Policy brief: Nurse fatigue, sleep, and health, and ensuring patient and public safety. *Nursing outlook*, 67(5), 615–619. <https://doi.org/10.1016/j.outlook.2019.08.004>

Cha, C., & Baek, G. (2023). Factors influencing the burnout dimensions among nurses: A cross-sectional study in South Korea. *Nursing open*, 10(12), 7725–7737. <https://doi.org/10.1002/nop2.2013>

Chen, Y. H., Yeh, C. J., Pan, L. F., & Jong, G. P. (2022). Relationships between Alcohol Use, Musculoskeletal Pain, and Work-Related Burnout. *Medicina (Kaunas, Lithuania)*, 58(8), 1022. <https://doi.org/10.3390/medicina58081022>

de Castro, A. B., Fujishiro, K., Rue, T., Tagalog, E. A., Samaco-Paquiz, L. P., & Gee, G. C. (2010). Associations between work schedule characteristics and occupational injury and illness. *International nursing review*, 57(2), 188–194. <https://doi.org/10.1111/j.1466-7657.2009.00793.x>

Dillman, D. A., Smyth, J. D., & Christian, L. M. (2014). *Internet, Phone, Mail, and Mixed-Mode Surveys: The Tailored Design Method* (4th ed.). John Wiley & Sons.

Dion, G., & Tessier, R. (1994). Validation de la traduction de l’Inventaire d’épuisement professionnel de Maslach et Jackson. *Canadian Journal of Behavioural Science / Revue canadienne des sciences du comportement*, 26(2), 210–227. <https://doi.org/10.1037/0008-400X.26.2.210>

Dupret, E., Bocérean, C., Teherani, M., Feltrin, M., & Pejtersen, J. H. (2012). Psychosocial risk assessment: French validation of the Copenhagen Psychosocial Questionnaire (COPSOQ). *Scandinavian journal of public health*, 40(5), 482–490. <https://doi.org/10.1177/1403494812453888>

Dyrbye, L., West, C., Johnson, P., Cipriano, P., Peterson, C., Beatty, D., Major-Elechi, B., & Shanafelt, T. (2020). Original Research: An Investigation of Career Choice Regret Among American Nurses. *The American journal of nursing*, 120(4), 24–33. <https://doi.org/10.1097/01.NAJ.0000660020.17156.ae>

Fédération interprofessionnelle de la santé du Québec. (2017). *Heures Supplémentaires. Agir collectivement, Agir localement*. https://affilies.fiqsante.qc.ca/cantons-de-lest/wp-content/uploads/sites/6/2020/10/TS_FIQ.pdf?download=1

Fernet, C., Trépanier, S. G., Demers, M., & Austin, S. (2017). Motivational pathways of occupational and organizational turnover intention among newly registered nurses in Canada. *Nursing outlook*, 65(4), 444–454. <https://doi.org/10.1016/j.outlook.2017.05.008>

Gao, F., Chui, P. L., Che, C. C., & Mao, X. R. (2023). Nurses’ Burnout, Resilience, and Its Associated Factors in Sichuan, China, During the Peak of Coronavirus Disease 2019 Infection. *Florence Nightingale journal of nursing*, 31(3), 152–159. <https://doi.org/10.5152/FNIN.2023.23023>

Geiger-Brown, J., Trinkoff, A., & Rogers, V. E. (2011). The impact of work schedules, home, and work demands on self-reported sleep in registered nurses. *Journal of occupational and environmental medicine*, 53(3), 303–307. <https://doi.org/10.1097/JOM.0b013e31820c3f87>

Government of Canada. (2024, June 3). *Hours of work - Federally regulated workplaces*. <https://www.canada.ca/en/services/jobs/workplace/federal-labour-standards/work-hours.html#h3.01.01>

IBM. (2019). *IBM SPSS Statistics for Windows* (version 26.0) [Computer software]. IBM Corp. <https://www.ibm.com/support/pages/downloading-ibm-spss-statistics-26-end-support-30-sep-2025>

International COPSOQ Network (2025). <http://www.copsoq-network.org>

Jacobson Consulting Inc. (2017). *Trends in Own Illness- or Disability-Related Absenteeism and Overtime Among Publicly Employed Registered Nurses. Quick Facts 2017*. Canadian Federation of Nurses Unions. https://nursesunions.ca/wp-content/uploads/2017/05/Quick_Facts_Absenteeism-and-Overtime-2017-Final.pdf

Kalisch, B. J., & Williams, R. A. (2009). Development and psychometric testing of a tool to measure missed nursing care. *The Journal of nursing administration*, 39(5), 211–219. <https://doi.org/10.1097/NNA.0b013e3181a23cf5>

Kunaviktikul, W., Wichaikhum, O., Nantsupawat, A., Nantsupawat, R., Chontawan, R., Klunklin, A., Roongruangsri, S., Nantachaipan, P., Supamanee, T., Chitpakdee, B., Akkadechanunt, T., & Sirakamon, S. (2015). Nurses' extended work hours: Patient, nurse and organizational outcomes. *International Nursing Review*, 62(3), 386–393. <https://doi.org/10.1111/inr.12195>

Lavoie-Tremblay, M., Trépanier, S. G., Fernet, C., & Bonneville-Roussy, A. (2014). Testing and extending the triple match principle in the nursing profession: a generational perspective on job demands, job resources and strain at work. *Journal of advanced nursing*, 70(2), 310–322. <https://doi.org/10.1111/jan.12188>

Lheureux, F., Truchot, D., Borteyrou, X., & Rascle, N. (2017). The Maslach Burnout Inventory – Human Services Survey (MBI-HSS): factor structure, wording effect and psychometric qualities of known problematic items. *Le travail humain*, 80(2), 161–186. <https://doi.org/10.3917/th.802.0161>

Low, G. K., Subedi, S., Omosumwen, O. F., Jhee, S. F., Devkota, S., Shanmuganathan, S., & Doyle, Z. (2024). Development and validation of observational and qualitative study protocol reporting checklists for novice researchers (ObsQual checklist). *Evaluation and program planning*, 106, 102468. <https://doi.org/10.1016/j.evalprogplan.2024.102468>

Medscape RN/LPN compensation report, 2020, October 28 [PowerPoint].

Min, A., Yoon, Y. S., Hong, H. C., & Kim, Y. M. (2019) Association between nurses' breaks, missed nursing care and patient safety in Korean hospitals. *Journal of Nursing Management*, 28(8), 2266–2274. <https://doi.org/10.1111/jonm.12831>

Naylor, J., Gillespie, G. L., Betcher, C., & Orr, C. E. (2022). Cost Analysis of Providing Overtime to Current Nurses Versus Hiring a Dedicated Nurse for COVID-19 Management in a Processing Plant. *Workplace health & safety*, 70(1), 24–30. <https://doi.org/10.1177/21650799211027868>

Nogues, S., & Tremblay, D. G. (2023). From Chosen to Forced: A Qualitative Exploration of Nurses' Experiences With Overtime. *Employee Responsibilities and Rights Journal*. <https://doi.org/10.1007/s10672-023-09485-y>

O'Driscoll, M. P., & Beehr, T. A. (1994). Supervisor behaviors, role stressors and uncertainty as predictors of personal outcomes for subordinates. *Journal of Organizational Behavior*, 15(2), 141–155. <https://doi.org/10.1002/job.4030150204>

Occelli, P., Quenon, J. L., Kret, M., Domecq, S., Delaperche, F., Claverie, O., Castets-Fontaine, B., Amalberti, R., Auroy, Y., Parneix, P., & Michel, P. (2013). Validation of the French version of the Hospital Survey on Patient Safety Culture questionnaire. *International journal for quality in health care: journal of the International Society for Quality in Health Care*, 25(4), 459–468. <https://doi.org/10.1093/intqhc/mzt047>

Palese, A., Navone, E., Danielis, M., Vryonides, S., Sermeus, W., & Papastavrou, E. (2021). Measurement tools used to assess unfinished nursing care: A systematic review of psychometric properties. *Journal of advanced nursing*, 77(2), 565–582. <https://doi.org/10.1111/jan.14603>

Patrician, P. A. (2004). Single-item graphic representational scales. *Nursing research*, 53(5), 347–352. <https://doi.org/10.1097/00006199-200409000-00011>

Sorra, J. S., & Dyer, N. (2010). Multilevel psychometric properties of the AHRQ hospital survey on patient safety culture. *BMC Health Services Research*, 10, 199. <https://doi.org/10.1186/1472-6963-10-199>

Statistics Canada. (2020, September 1). *Overtime work among professional nurses during the COVID-19 pandemic*. <https://www150.statcan.gc.ca/n1/pub/45-28-0001/2020001/article/00074-eng.htm>

Statistics Canada. (2023, July 24). *Nurses: Working harder, more hours amid increased labour shortage*. <https://www.statcan.gc.ca/01/en/plus/4165-nurses-working-harder-more-hours-amid-increased-labour-shortage>

Stone, P. W., Mooney-Kane, C., Larson, E. L., Horan, T., Glance, L. G., Zwanziger, J., & Dick, A. W. (2007). Nurse working conditions and patient safety outcomes. *Medical care*, 45(6), 571–578. <https://doi.org/10.1097/MLR.0b013e3180383667>

Tabachnick, B. G., Fidell, L. S. (2019). *Using multivariate statistics* (7th ed.). Pearson.

Trinkoff, A. M., Johantgen, M., Storr, C. L., Gurses, A. P., Liang, Y., & Han, K. (2011). Nurses' work schedule characteristics, nurse staffing, and patient mortality. *Nursing research*, 60(1), 1–8. <https://doi.org/10.1097/NNR.0b013e3181fff15d>

Vallerand R. J., & Halliwell W. R. (1983). Vers une méthodologie de validation culturelle de questionnaires psychologiques : implications pour la psychologie du sport [Toward a methodology of cross-cultural validation of psychological questionnaire : implications for psychology of sport]. *Canadian Journal of Applied Sport Sciences* 8, 9–18.

von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gøtzsche, P. C., Vandenbroucke, J. P., & STROBE Initiative (2007). The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: Guidelines for reporting observational studies. *Lancet*, 370(9596), 1453–1457. [https://doi.org/10.1016/S0140-6736\(07\)61602-X](https://doi.org/10.1016/S0140-6736(07)61602-X)

Watanabe, M., & Yamauchi, K. (2016). Psychosocial Factors of Overtime Work in Relation to Work-Nonwork Balance: a Multilevel Structural Equation Modeling Analysis of Nurses Working in Hospitals. *International journal of behavioral medicine*, 23(4), 492–500. <https://doi.org/10.1007/s12529-016-9563-x>

Watanabe, M., & Yamauchi, K. (2018). The effect of quality of overtime work on nurses' mental health and work engagement. *Journal of nursing management*, 26(6), 679–688. <https://doi.org/10.1111/jonm.12595>

Watanabe, M., & Yamauchi, K. (2019). Subtypes of overtime work and nurses' fatigue, mental status, and work engagement: A latent class analysis of Japanese hospital nurses. *Journal of advanced nursing*, 75(10), 2122–2132. <https://doi.org/10.1111/jan.13991>

White, E. M., Aiken, L. H., & McHugh, M. D. (2019). Registered Nurse Burnout, Job Dissatisfaction, and Missed Care in Nursing Homes. *Journal of the American Geriatrics Society*, 67(10), 2065–2071. <https://doi.org/10.1111/jgs.16051>

Winwood, P. C., Winefield, A. H., Dawson, D., & Lushington, K. (2005). Development and validation of a scale to measure work-related fatigue and recovery: the Occupational Fatigue Exhaustion/Recovery Scale (OFER). *Journal of occupational and environmental medicine*, 47(6), 594–606. <https://doi.org/10.1097/01.jom.0000161740.71049.c4>

Winwood, P. C., Lushington, K., & Winefield, A. H. (2006). Further development and validation of the Occupational Fatigue Exhaustion Recovery (OFER) scale. *Journal of occupational and environmental medicine*, 48(4), 381–389. <https://doi.org/10.1097/01.jom.0000194164.14081.06>

Appendix 1

Table 1

Project Timeline

Activity	Dates
• The development of study design	June-July 2024
• Ethics approval	November 2024
• Translation and adaptation in the French of MISSCARE Survey instrument	December 2024-May 2025
• Cross-sectional study: Data collection	May-June 2025
• Cross-sectional study: Analysis	July-August 2025
• Cross-section study: Submission of manuscripts and communications	September 2025-February 2026